

Literaturverzeichnis

zum Titelthema „S3-Leitlinie - Prävention und Therapie der Adipositas“ von Professor Dr. Hans Hauner, Bayerisches Ärzteblatt 7-8/2016, Seite 344 ff.

1. Kurth BM. Erste Ergebnisse aus der „Studie zur Gesundheit Erwachsener in Deutschland“ (DEGS) . Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2012;55(8):980-90.
2. Blüher S, Meigen C, Gausche R, Keller E, Pfäffle R, Sabin M et al.: Age-specific stabilization in obesity prevalence in German children: a cross-sectional study from 1999 to 2008. Int J Pediatr Obes 2011;6:e199-206.
3. World Health Organization. Obesity – preventing and managing the global epidemic. Report of a WHO Consultation on obesity. Technical Report Series 894, 2000, Geneva
4. Hauner H. Secretory factors from human adipose tissue and their functional role. Proc Nutr Soc 2005;64:163-9
5. Knoll K-P, Hauner H: Kosten der Adipositas in der Bundesrepublik Deutschland. Eine aktuelle Krankheitskostenstudie. Adipositas 2008;2:204-10.
6. Effertz T, Engel S, Verheyen F, Linder R. The costs and consequences of obesity in Germany: a new approach from a prevalence and life-cycle perspective. Eur J Health Econ 23 December 2015.
7. Wirth A, Wabitsch M, Hauner H. Klinische Leitlinie: Prävention und Therapie der Adipositas. Dtsch Ärztebl 2014;111:705-13.
8. Hauner H, Moss A, Barg A, Bischoff SC, Colombo-Benkmann M, Ellrott T et al. Interdisziplinäre Leitlinie der Qualität S3 zur “Prävention und Therapie der Adipositas”. Adipositas 2014;8:179-221.
9. Canoy D, Boekholdt SM, Wareham N, Luben R, Welch A, Bingham S et al. Body fat distribution and risk of coronary heart disease in men and women in the European Prospective Investigation Into Cancer and Nutrition in Norfolk cohort: a population-based prospective study. Circulation 2007;116(25):2933-43.
10. Prospective Studies Collaboration. Body-mass index and cause-specific mortality in 900 000 adults. Collaborative analyses of 57 prospective studies. Lancet 2009;373:1083-96.
11. De Zwaan M., Petersen I, Kaerber M, Burgmer R, Nolting B, Legenbauer T et al. Obesity and quality of life: a controlled study of normal-weight and obese individuals. Psychosomatics 2009;50(5):474-82.
12. Hilbert A, Rief W, Braehler E. Stigmatizing attitudes toward obesity in a representative population-based sample. Obesity (Silver Spring) 2008;16(7):1529-34.
13. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, Zitman FG. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. Arch Gen Psychiatry 2010;67(3):220-9.
14. Bes-Rastrollo M, van Dam RM, Martinez-Gonzalez MA, Li TY, Sampson LL, Hu FB. Prospective study of dietary energy density and weight gain in women. Am J Clin Nutr 2008;88(3):769-77.
15. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. Med Sci Sports Exerc 2009;41(2):459-71.
16. Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. N Engl J Med 2008;359(3):229-4.

17. Sacks FM, Bray GA, Carey VJ, Smith SR, Ryan DH, Anton SD et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med* 2009;360(9):859-73.
18. Johnston BC, Kanters S, Bandayrel K, Wu P, Naji F, Siemienjuk RA et al. Comparison of weight loss among named diet programs in overweight and obese adults. A meta-analysis. *JAMA* 2014;312:923-33.
19. Anderson JW, Reynolds LR, Bush HM, Rinsky JL, Washnock C. Effect of a behavioral/nutritional intervention program on weight loss in obese adults: a randomized controlled trial. *Postgrad Med* 2011;123(5):205-13.
20. Ohsiek S, Williams M. Psychological factors influencing weight loss maintenance: an integrative literature review. *J Am Acad Nurse Pract* 2011;23(11):592-601.
21. Toubro S, Astrup A. Randomised comparison of diets for maintaining obese subjects' weight after major weight loss: ad lib, low fat, high carbohydrate diet v fixed energy intake. *BMJ* 1997;314(7073):29-34.
22. Butryn ML, Phelan S, Hill JO, Wing RR. Consistent self-monitoring of weight: a key component of successful weight loss maintenance. *Obesity (Silver Spring)* 2007;15(12):3091-6.
23. Monami M, Dicembrini I, Marchionni N, Rotella CM, Mannucci E. Effects of glucagon-like Peptide-1 receptor agonists on body weight: a meta-analysis. *Exp Diabetes Res* 2012;2012:672658.
24. Sjöström L, Narbro K, Sjöström CD, Karason K, Larsson B, Wedel H et al.: Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med* 2007;357:741-52
25. Sjöström L Review of the key results from the Swedish obese subjects (SOS) trial – a prospective controlled intervention study of bariatric surgery. *J Intern Med* 2013;373:219-34.
26. Schauer PR, Bhatt DL, Kirwan JP, Wolski K, Brethauer SA, Navaneethan SD et al. Bariatric surgery versus intensive medical therapy for diabetes – 3-year outcomes. *N Engl J Med* 2014;370(21):2002-13.

Statistisches Bundesamt. Mikrozensus - Fragen zur Gesundheit - Körpermaße der Bevölkerung. 2011 [cited: 2013 Mai 30].