

Literaturverzeichnis zum Titelthema “Neues aus der Neurochirurgie”

Bayerisches Ärzteblatt 10/2009, Seite 464 ff.

Dr. Karl-Michael Schebesch und Dr. Jan Warnat

- 1 Localized BCNU chemotherapy and the multimodal management of malignant glioma.
La Rocca RV, Mehdorn HM.
Curr Med Res Opin. 2009 Jan;25(1):149-60.
- 2 Awake craniotomy for brain tumors near eloquent cortex: correlation of intraoperative cortical mapping with neurological outcomes in 309 consecutive patients.
Kim SS, McCutcheon IE, Suki D, Weinberg JS, Sawaya R, Lang FF, Ferson D, Heimberger AB, DeMonte F, Prabhu SS.
Neurosurgery. 2009 May;64(5):836-45; discussion 345-6.
- 3 Preoperative and intraoperative diffusion tensor imaging-based fiber tracking in glioma surgery.
Nimsky C, Ganslandt O, Hastreiter P, Wang R, Benner T, Sorensen AG, Fahlbusch R.
Neurosurgery. 2007 Jul;61(1 Suppl):178-85; discussion 186.
- 4 A multinational comparison of subarachnoid hemorrhage epidemiology in the WHO MONICA stroke study.
Ingall T, Asplund K, Mähönen M, Bonita R.
Stroke. 2000 May;31(5):1054-61.
- 5 Subarachnoid haemorrhage: diagnosis, causes and management.
van Gijn J, Rinkel GJ.[^] *Brain.* 2001 Feb;124(Pt 2):249-78. Review.
- 6 Systemic treatments of the vasospasm
Tavernier B, Decamps F, Vega E, Poidevin P, Verdin M, Riegel B
Ann Fr Anesth Reanim. 2007 Nov;26(11):980-4.
- 7 Clazosentan to overcome neurological ischemia and infarction occurring after subarachnoid hemorrhage (CONSCIOUS-1): randomized, double-blind, placebo-controlled phase 2 dose-finding trial.
Macdonald RL, Kassell NF, Mayer S, Ruefenacht D, Schmiedek P, Weidauer S, Frey A, Roux S, Pasqualin A; CONSCIOUS-1 Investigators.
Stroke. 2008 Nov;39(11):3015-21. Epub 2008 Aug 7.
- 8 International Subarachnoid Aneurysm Trial (ISAT) of neurosurgical clipping versus endovascular coiling in 2143 patients with ruptured intracranial aneurysms: a randomised trial.
Molyneux A, Kerr R, Stratton I, Sandercock P, Clarke M, Shrimpton J, Holman R; International Subarachnoid Aneurysm Trial (ISAT) Collaborative Group.
Lancet. 2002 Oct 26;360(9342):1267-74.
- 9 Unruptured intracranial aneurysms: natural history, clinical outcome, and risks of surgical and endovascular treatment.
Wiebers DO, Whisnant JP, Huston J 3rd, Meissner I, Brown RD Jr, Piepgras DG, Forbes GS, Thielen K, Nichols D, O'Fallon WM, Peacock J, Jaeger L, Kassell NF, Kongable-Beckman GL, Torner JC; International Study of Unruptured Intracranial Aneurysms Investigators.
Lancet. 2003 Jul 12;362(9378):103-10.
- 10 Microsurgical management of incompletely coiled and recurrent aneurysms: trends, techniques, and observations on coil extrusion.

- Waldron JS, Halbach VV, Lawton MT.
Neurosurgery. 2009 May;64(5 Suppl 2):301-15; discussion 315-7.
- 11 Failure of extracranial-intracranial arterial bypass to reduce the risk of ischemic stroke.
Results of an international randomized trial. The EC/IC Bypass Study Group.
[No authors listed]
N Engl J Med. 1985 Nov 7;313(19):1191-200.
- 12 Clinical outcome of standard extracranial-intracranial bypass surgery in patients with symptomatic atherosclerotic occlusion of the internal carotid artery.
Mendelowitsch A, Taussky P, Rem JA, Gratzl O.
Acta Neurochir (Wien). 2004 Feb;146(2):95-101. Epub 2004 Jan 30.
- 13 Deep hypothermic circulatory arrest for complex cerebral aneurysms: lessons learned.
Mack WJ, Ducruet AF, Angevine PD, Komotar RJ, Shrebnick DB, Edwards NM, Smith CR, Heyer EJ, Monyero L, Connolly ES Jr, Solomon RA.
Neurosurgery. 2008 Jun;62(6 Suppl 3):1311-23.
- 14 Mummaneni PV, Burkus JK, Haid RW, Traynelis VC, Zdeblick TA.
Clinical and radiographic analysis of cervical disc arthroplasty compared with allograft fusion: a randomized controlled clinical trial.
J Neurosurg Spine. 2007 Mar;6(3):198-209.
- 15 Chang UK, Kim DH, Lee MC, Willenberg R, Kim SH, Lim J. Range of motion change after cervical arthroplasty with ProDisc-C and prestige artificial discs compared with anterior cervical discectomy and fusion.
J Neurosurg Spine. 2007 Jul;7(1):40-6.
- 16 Heller JG, Sasso RC, Papadopoulos SM, Anderson PA, Fessler RG, Hacker RJ, Coric D, Cauthen JC, Riew DK. Comparison of BRYAN cervical disc arthroplasty with anterior cervical decompression and fusion: clinical and radiographic results of a randomized, controlled, clinical trial.
Spine. 2009 Jan 15;34(2):101-7.
- 17 Sasso RC, Smucker JD, Hacker RJ, Heller JG.
Artificial disc versus fusion: a prospective, randomized study with 2-year follow-up on 99 patients. Spine. 2007 Dec 15;32(26):2933-40.
- 18 Guyer RD, McAfee PC, Banco RJ, Bitan FD, Cappuccino A, Geisler FH, Hochschuler SH, Holt RT, Jenis LG, Majd ME, Regan JJ, Tromanhauser SG, Wong DC, Blumenthal SL. Prospective, randomized, multicenter Food and Drug Administration investigational device exemption study of lumbar total disc replacement with the CHARITE artificial disc versus lumbar fusion: five-year follow-up.
Spine J. 2009 May;9(5):374-86.
- 19 Cakir B, Carazzo C, Schmidt R, Mattes T, Reichel H, Käfer W.
Adjacent segment mobility after rigid and semirigid instrumentation of the lumbar spine. Spine. 2009 May 20;34(12):1287-91.
- 20 Deuschl G, Schade-Brittinger C, Krack P, Volkmann J, Schäfer H, Bötzl K, Daniels C, Deutschländer A, Dillmann U, Eisner W, Gruber D, Hamel W, Herzog J, Hilker R, Klebe S, Kloss M, Koy J, Krause M, Kupsch A, Lorenz D, Lorenzl S, Mehdorn HM, Moringlane JR, Oertel W, Pinsker MO, Reichmann H, Reuss A, Schneider GH, Schnitzler A, Steude U, Sturm V, Timmermann L, Tronnier V, Trottenberg T, Wojtecki L, Wolf E, Poewe W, Voges J; German Parkinson Study Group, Neurostimulation Section.

- A randomized trial of deep-brain stimulation for Parkinson's disease.
N Engl J Med. 2006 Aug 31;355(9):896-908.
- 21 Weaver FM, Follett K, Stern M, Hur K, Harris C, Marks WJ Jr, Rothlind J, Sagher O, Reda D, Moy CS, Pahwa R, Burchiel K, Hogarth P, Lai EC, Duda JE, Holloway K, Samii A, Horn S, Bronstein J, Stoner G, Heemskerk J, Huang GD; CSP 468 Study Group. Bilateral deep brain stimulation vs best medical therapy for patients with advanced Parkinson disease: a randomized controlled trial. JAMA. 2009 Jan 7;301(1):63-73.
- 22 Vidailhet M, Vercueil L, Houeto JL, Krystkowiak P, Lagrange C, Yelnik J, Bardinet E, Benabid AL, Navarro S, Dormont D, Grand S, Blond S, Ardouin C, Pillon B, Dujardin K, Hahn-Barma V, Agid Y, Destée A, Pollak P; French SPIDY Study Group. Bilateral, pallidal, deep-brain stimulation in primary generalised dystonia: a prospective 3 year follow-up study. Lancet Neurol. 2007 Mar;6(3):223-9.
- 23 Schrader C, Benecke R, Deuschl G, Hilker R, Kupsch A, Lange M, Sixel-Döring F, Timmermann L, Volkmann J, Fogel W. Deep brain stimulation for dystonia: Consensus recommendations of the German Deep Brain Stimulation Association. Nervenarzt. 2009 Apr 30.
- 24 Timmermann L, Deuschl G, Fogel W, Hilker R, Kupsch A, Lange M, Schrader C, Sixel-Döring F, Volkmann J, Benecke R. Deep brain stimulation for tremor in multiple sclerosis: Consensus recommendations of the Deep Brain Stimulation Association. Nervenarzt. 2009 May 27.
- 25 Sixel-Döring F, Benecke R, Fogel W, Hilker R, Kupsch A, Lange M, Schrader C, Timmermann L, Volkmann J, Deuschl G. Deep brain stimulation for essential tremor: Consensus recommendations of the German Deep Brain Stimulation Association. Nervenarzt. 2009 Apr 30.
- 26 Hilker R, Benecke R, Deuschl G, Fogel W, Kupsch A, Schrader C, Sixel-Döring F, Timmermann L, Volkmann J, Lange M. Deep brain stimulation for Parkinson's disease: Consensus recommendations of the German Deep Brain Stimulation Association. Nervenarzt. 2009 Apr 11.