

Literaturverzeichnis
zum Titelthema „Diagnostik und Therapie der Venenthrombose und der Lungenembolie“
Bayerisches Ärzteblatt 5/2015, Seite 216 ff.
von Dr. Thomas Noppene

1. Akl EA, Barba M, Rohilla S et al. Low-molecular-weight heparins are superior to vitamin K antagonists for the long term treatment of venous thromboembolism in patients with cancer: a cochrane systematic review.
J Exp Clin Cancer Res 2008; 27: Seiten 21
2. Cohen AT, Agnelli G, Anderson FA, Arcelus JI et al. Venous thromboembolism (VTE) in Europe. The number of VTE events and associated morbidity and mortality.
Thromb Haemost 2007; 98:756-764
3. Flinterman LE, van Hylckama Vlieg A, Cannegieter SC et al. Long-Term Survival in a large Cohort of Patients with Venous Thrombosis: Incidence and Predictors.
PLoS Med (2012); 9:e10001155.
4. Hach-Wunderle V, Blättler W, Gerlach H et al. Diagnostik und Therapie der Venenthrombose und der Lungenembolie. Interdisziplinäre S2 Leitlinie.
VASA 2010 (Suppl.): S78
5. Hirsh J, Dalen J, Anderson DR et al. Oral anticoagulants: mechanism of action, clinical effectiveness, and optimal therapeutic range.
Chest 2001; 119:8S-21S
6. Kahn SR, Shapiro S, Wells PS et al. Compression stockings to prevent post-thrombotic syndrome: a randomised placebo-controlled trial
Lancet. 2014; 383: 880-888
7. Kahn SR, Shapiro S, Ducruet T et al. Graduated compression stockings to treat acute leg pain associated with proximal DVT. A randomized controlled trial.
Thromb Haemost. 2014; 112: 1137-41.
8. Kearon C, Akl. E, Comerota AJ, et al. Antithrombotic Therapy for VTE Disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed:
American College of Chest Physicians, Evidence-Based Clinical Practice Guidelines.
Chest 2012; 141:e419S-e494S
9. Lindhoff-Last E, Luxembourg B: Evidence-based indications for thrombophilia screening.
Vasa 2008; 37:19-30
10. Perkins JM, Magee TR, Galland RB: Phlegmasia caerulea dolens and venous gangrene.
Br J Surg 1996;83:19-23
11. Schulman S, Rhedin AS, Lindmarker P, et al. A comparison of six weeks with six months of oral anticoagulant therapy after a first episode of venous thromboembolism. Duration of Anticoagulation Trial Study Group
N Engl J Med. 1995 Jun 22; 332 (25): 1661-5
12. Spyropoulos AC: Upper vs. lower extremity deep vein thrombosis: Outcome definition of venous thromboembolism for clinical predictor rules or risk factor analyses in hospitalized patients.
J Thromb Haemost 2009;7:1041-1042

13. Warkentin TE, Cook DJ: Current perspective of venous thrombosis in the upper extremity: a rebuttal. *J Thromb Haemost* 2009;7:237-238
14. Wells PS, Anderson DR, Rodger M et. al. Evaluation of D-dimer in the diagnosis of suspected deep-vein thrombosis. *N Engl J Med* 2003;349:1227-1235