

Literatur zum Varia-Beitrag "Früherkennung peripartaler Depressionen – Von der UPlusE-Studie in die Regelversorgung"

Bayerisches Ärzteblatt 9/2026, Seite 370 ff.

Autorinnen: Carolin Seivert und Dr. Susanne Simen

1. Liu X, Wang S, Wang G. Prevalence and risk factors of postpartum depression in women: a systematic review and meta-analysis. *J Clin Nurs*. 2021. doi:10.1111/jocn.16121.
2. Cameron EE, Sedov ID, Tomfohr-Madsen LM. Prevalence of paternal depression in pregnancy and the postpartum: an updated meta-analysis. *J Affect Disord*. 2016;206:189-203. doi:10.1016/j.jad.2016.07.044.
3. Gidén K, Vinnerljung L, Iliadis SI, Fransson E, Skalkidou A. Feeling better? Identification, interventions, and remission among women with early postpartum depressive symptoms in Sweden: a nested cohort study. *Eur Psychiatry*. 2024;67:e6. doi:10.1192/j.eurpsy.2024.6.
4. Place JMS, Renbarger K, Van De Griend K, Guinn M, Wheatley C, Holmes O. Barriers to help-seeking for postpartum depression mapped onto the socio-ecological model and recommendations to address barriers. *Front Glob Womens Health*. 2024;5:1335437. doi:10.3389/fgwh.2024.1335437.
5. Dorsch VM, Rohde A. Postpartale psychische Störungen—Update 2016. *Frauenheilkd Up2date*. 2016;10(4):355-374.
6. Henshaw EJ, Cooper M, Wood T, Doan SN, Krishna S, Lockhart M. Psychosocial predictors of early postpartum depressive and anxious symptoms in primiparous women and their partners. *BMC Pregnancy Childbirth*. 2023;23:55. doi:10.1186/s12884-023-05506-8.
7. Glasser S, Lerner-Geva L. Focus on fathers: paternal depression in the perinatal period. *Perspect Public Health*. 2019;139(4):195-198. doi:10.1177/1757913918790597.
8. Stein A, Pearson RM, Goodman SH, Rapa E, Rahman A, McCallum M, Howard LM, Pariante CM. Effects of perinatal mental disorders on the fetus and child. *Lancet*. 2014;384(9956):1800-1819. doi:10.1016/S0140-6736(14)61277-0.
9. COST Action Research Innovation and Sustainable Pan-European Network in Peripartum Depression Disorder (RISEUP-PPD). Evidence-based clinical practice guidelines for prevention, screening and treatment of peripartum depression [Internet]. 2025 [cited 2026 Jul 27]. Available from: <https://riseupppd18138.com/clinical-practice-guidelines-html/>
10. Zimmerman D, Verbov G, Edelstein N, Stein-Zamir C. Preventive health services for young children in Israel: historical development and current challenges. *Isr J Health Policy Res*. 2019;8:21. doi:10.1186/s13584-019-0287-7.
11. Royal College of Psychiatrists. Perinatal mental health services: recommendations for the provision of services for childbearing women. College Report CR232 [Internet]. London: Royal College of Psychiatrists; 2021 [cited 2026 Jul 27]. Available from: <https://www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/college-reports/college-report-cr232---perinatal-mental-health-services.pdf>
12. Stentzel U, et al. Identification and transfer to stepped care of depressed and psychosocially stressed parents during peri- and postpartum—UPlusE: study protocol for cluster randomized trial of a screening intervention. *Trials*. 2024;25:610.
13. Wilcox M, McGee B, Ionescu D, Leonte M, LaCross L, Reps J, Wildenhaus K. Perinatal depressive symptoms often start in the prenatal rather than postpartum period: results from a longitudinal study. *Arch Womens Ment Health*. 2021;24:119-131. doi:10.1007/s00737-020-01017-z.
14. Putnick DL, Sundaram R, Bell EM, Ghassabian A, Goldstein RB, Robinson SL, Vafai Y, Gilman SE, Yeung EH. Trajectories of maternal postpartum depressive symptoms. *Pediatrics*. 2020;146(5):e20200857. doi:10.1542/peds.2020-0857.